



FOR COUPLES WHERE ONE OF YOU HAS ADHD

# THE RSD CYCLE MAP

Rejection-sensitive dysphoria (RSD) does not run as a solo problem. When one partner has ADHD, RSD plays out between two people, and the cycle can run on its own regardless of what either of you wants. This is a shared vocabulary so you can spot it together.

## THREE THINGS TO KNOW FIRST

### <1s

#### FAST

The pain of RSD usually arrives in under a second, before conscious thought has time to catch up, which is why cognitive strategies like 'just think about it differently' do not land in the moment when the wave is already breaking.

### 0

#### NOT CHOSEN

This is not weaponized sensitivity or a character flaw — it is a nervous-system response common in ADHD that both partners can learn to see, name, and hold together instead of taking personally.

### 2

#### BETWEEN YOU

RSD plays out more between two people than it does inside one, so naming the cycle as ours (rather than yours) is what shifts it from a fight about who is the problem to a shared attention on the pattern itself.

## THE CYCLE IN FOUR PHASES

The cycle usually runs in this order. Read it together, name where you tend to enter it, and where you tend to exit.

### 1

#### THE TRIGGER

A small comment, a pause, a pointed silence, or a tone shift lands sideways — often something a neurotypical brain would barely register, but that hits the ADHD nervous system like a full alarm.

### 2

#### THE RESPONSE

The ADHD partner's system floods, and the reaction comes out disproportionate to what just happened, which usually surprises both partners once the wave has passed.

### 3

#### THE EDIT

The non-ADHD partner starts being careful, softening tone, avoiding topics, and walking on eggshells to keep the peace and to try to prevent the next flood.

### 4

#### THE CONFIRMATION

The ADHD partner senses the carefulness and reads it as proof that the original fear was accurate ('I really am too much'), and the fear then becomes the lived experience of the relationship.

### READ THIS TOGETHER, OUT LOUD

*"A comment lands sideways, and my system floods before I have a chance to think it through. You feel the wave and start being careful, and your carefulness lands as proof that I really am too much. Neither of us chose this, both of us maintain it without meaning to, and we are on the same side against the cycle rather than against each other."*

## HOW THE CYCLE SOUNDS IN REAL LIFE

### A COMPOSITE SCENARIO

You are three months into a pattern where every casual comment about the dishes has been landing like a personal indictment, and here is what one Tuesday evening looks like inside your own head.

**Partner:** “Did you forget to run the dishwasher?”

Your chest hits before your brain has any chance to catch up, and underneath the question you hear something more like “you are incompetent, and I am losing patience with you,” even though nobody actually said that. Within ten seconds you have escalated to “I can never do anything right around you” and you have already left the room, while your partner is standing in the kitchen having asked a five-word question about a dishwasher. This cycle has run some version of itself thirty times before.

This time, you come back after twenty-five minutes and try the re-approach the two of you agreed to when things were calm.

**You:** “I was in a wave. Can you tell me what you actually meant?”

**Partner:** “I meant if you didn't run it, I'll run it now.”

**Your body finally lets go of the story that had been running for months. The cycle itself did not stop existing that night, but this one time it did not get to run the relationship — and over enough repetitions, that adds up to the cycle keeping existing but no longer shaping who the two of you are to each other.**

## SET UP THE PAUSE BEFORE THE NEXT WAVE

The pause tool only works when it is pre-agreed. Trying to invent it in the middle of a wave is exactly how couples lose it before it can help. Set these four things when you are both regulated.

### SIGNAL WORD

Choose one phrase either of you can say without explanation, and pre-agree that saying it means we pause immediately, without debating in the moment whether the RSD is 'actually' happening or whether the topic 'really' warrants a break.

*“I'm in a wave.” · “Pause please.” · “Red flag.” · A nonverbal signal you both know and use the same way.*

### WHERE YOU GO

Take a nervous-system break in separate physical space, ideally somewhere that actively changes your body state (movement, water, cold air, music), rather than staying in the same room and calling it a pause while the tension quietly stays in the air.

*A walk around the block. Different room with the door shut. Hot shower. Coffee run. Headphones on with music.*

### TIME BOUNDARY

Set a timer for 20 to 30 minutes minimum, come back when it goes off even if things still feel edgy, and if you need more time say so out loud rather than silently extending — an unannounced extension lands as abandonment and often starts the cycle over from scratch.

*“Timer is set for 25 minutes, and I'll be back then.” · “I need another 15 — is that OK?”*

### HOW YOU COME BACK

Use a pre-agreed soft phrase to signal you are re-approaching, and treat that phrase as the invitation to reconnect rather than as an opening line for re-litigating whatever triggered the wave in the first place.

*“I'm ready to try again.” · “I'm back, can we?” · “Let's re-approach when you have a minute.”*

## THE RE-APPROACH AFTER THE PAUSE

Coming back is a specific three-step re-entry, not a re-litigation of what triggered the wave. Each partner has a role.

### 1 NAME THE WAVE, NOT THE TRIGGER

Saying 'I was in an RSD wave' is more honest and more useful than 'you shouldn't have said that,' because the wave is the thing to name first — the trigger itself can be discussed once both nervous systems have actually settled.

#### PARTNER WITH ADHD MIGHT SAY

*"That reaction was bigger than the thing that started it, and I think that was an RSD wave, not a response to what you actually said."*

#### NON-ADHD PARTNER MIGHT SAY

*"I saw you get flooded, and I'm not going to try to argue the content while you're still coming back — are you back with me yet?"*

### 2 ASK WHAT WAS MEANT

RSD tends to supply a meaning that was not actually there, so checking what your partner meant out loud is what dissolves the confirmation phase before it hardens into a story the relationship has to live with.

#### PARTNER WITH ADHD MIGHT SAY

*"When you said X, what I heard was Y, and I want to check that against what you actually meant before I go any further with it."*

#### NON-ADHD PARTNER MIGHT SAY

*"What I actually meant was Z, which is different from what you heard — does that land differently now that I've said it plainly?"*

### 3 REPAIR THE EDIT

The non-ADHD partner naming the eggshells out loud is what closes the loop, because otherwise the ADHD partner is left to infer the eggshells alone, and inferring silence is exactly what RSD does worst.

#### PARTNER WITH ADHD MIGHT SAY

*"Thank you for naming that you were walking on eggshells — not knowing whether I was imagining it was making it worse in my head."*

#### NON-ADHD PARTNER MIGHT SAY

*"I've noticed I've been softening around this topic all week, and I want to name it so you're not left to infer it and fill in the reason yourself."*

## WHAT EACH PARTNER BRINGS BACK

The cycle changes when each partner has one job. Not fixing the other person — noticing your own move in the cycle and doing it differently.

#### THE PARTNER WITH ADHD

- Notice the wave as it starts, because even a small amount of space between the wave arriving and the reaction leaving your mouth is enough to change what happens next.
- Practice saying "I am in a wave" out loud without waiting for the reaction to prove it first, so the language becomes automatic before the escalation does.
- When the wave is already breaking and the pause is not yet available, DBT's TIPP skills (cold water on the face, thirty seconds of intense movement, or paced breathing with an exhale twice as long as the inhale) can shorten the physiological wave itself and buy you space.
- Give yourself grief and self-compassion for the years of waves that ran before either of you had any framework for what was happening. Living inside a pattern you could not describe produced its own loss, and processing that loss belongs alongside building the practical protocol, not after it.

#### THE NON-ADHD PARTNER

- Notice the edit — the specific moment you catch yourself softening tone past what feels honest, or avoiding a topic to keep the peace — so you can name it before it accumulates into a pattern.
- When you see the pattern take hold, name it as RSD (not as your partner's character), because the same behavior in those two different frames puts the two of you in two very different rooms.
- Say the pre-agreed phrase early rather than waiting until you are already frustrated, because frustration is exactly the state in which most couples lose the pause tool right when they need it.

**The cycle will run again**, and that is not evidence that the work isn't working — it's evidence that the pattern is well-practiced, and unlearning it takes reps rather than a single decision. Even one instance of naming the cycle out loud together while it is happening changes how the cycle can hold you afterward, because the pattern loses some of its grip the moment it stops being invisible. Some couples find that the wave itself softens with certain ADHD medications, most often guanfacine or clonidine per William Dadds's clinical work — that is a conversation for your